

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Occupational Exposure to Endocrine-Disrupting Chemicals and Birth Weight and Length of Gestation: A European Meta-Analysis

Laura Birks, Maribel Casas, Ana M. Garcia, Jan Alexander, Henrique Barros, Anna Bergström, Jens Peter Bonde, Alex Burdorf, Nathalie Costet, Asta Danileviciute, Merete Eggesbø, Mariana F. Fernández, M. Carmen González-Galarzo, Regina Gražulevičienė, Wojciech Hanke, Vincent Jaddoe, Manolis Kogevinas, Inger Kull, Aitana Lertxundi, Vasiliki Melaki, Anne-Marie Nybo Andersen, Nicolás Olea, Kinga Polanska, Franca Rusconi, Loreto Santa-Marina, Ana Cristina Santos, Tanja Vrijkotte, Daniela Zugna, Mark Nieuwenhuijsen, Sylvaine Cordier, and Martine Vrijheid

Table of Contents

Excel File Table S1: See “Additional Files” below.

Excel File Table S2: See “Additional Files” below.

Table S3: Coding of maternal education level by cohorts

Table S4: Distribution of covariates by cohorts

Table S5: Maternal occupational exposures to EDC groups during pregnancy as classified by a job exposure matrix and meta-analyzed associations (95%CI) with birth term LBW and length of gestation^a, excluding DNBC and MoBa cohorts

Table S6: Maternal occupational exposures to EDC groups during pregnancy as classified by a job exposure matrix and meta-analyzed associations (95%CI) with birth term LBW and length of gestation^a, excluding elected cesarean births

Table S7: Minimally adjusted models for maternal exposure to endocrine disrupting chemicals as classified by a job exposure matrix and associations with

Figure S1: Meta-analysis of odds ratios for term low birth weight for pregnant women occupationally exposed to 4 or more endocrine-disrupting chemical groups as classified by a job exposure matrix. N's

represent subjects included in complete case analysis. Cohorts excluded had no cases of term LBW among pregnant women classified as occupationally exposed to 4 or more endocrine-disrupting chemical groups. All models are adjusted for maternal age, parity, maternal education, maternal smoking, maternal BMI, marital status, sex of newborn, and race and gestational age, where applicable. Unexposed mothers are used as reference group. Shaded boxes around the point estimates indicate the weight of the study-specific estimate.

Figure S2: Meta-analysis of odds ratios for term LBW for pregnant women occupationally exposed to phthalates as classified by a job exposure matrix. N's represent subjects included in complete case analysis. Cohorts excluded had no cases of term LBW among pregnant women classified as occupationally exposed to 4 or more endocrine-disrupting chemical groups, except in Generation XXI, which had one case. All models are adjusted for maternal age, parity, maternal education, maternal smoking, maternal BMI, marital status, sex of newborn, and race and gestational age, where applicable. Unexposed mothers are used as reference group. Shaded boxes around the point estimates indicate the weight of the study-specific estimate.

Additional Files

Supplemental Code and Data Zip File

Supplemental Code and Data Zip File Index

Excel File Table S1: Application of Brouwers 2009 JEM to ISCO88 and SOC2000 codes using CAMSIS translation with exposure score

Excel File Table S2: Experts' assignment of proxy codes for use with the Brouwers 2009 JEM